Table 2: Volume Perfusion Computed Tomography perfusion parameters of Case 2

Parameters	ROI-1 (Pancreatic body mass)	ROI-2 (Pancreatic body mass)	ROI-3 (Normal pancreas)
PEI (HU)	57.05	63.32	110.33
BF (ml/100 gm/min)	11.59	8.12	51.68
BV (ml/100 gm)	2.82	3.07	20.76

encasing the celiac axis and its branches in the pancreatic body region. There was peri-pancreatic fat stranding and thickening of renal fascia with a contiguous subdiaphragmatic collection on left side. There was no associated retroperitoneal or mesenteric lymphadenopathy, thickened bowel loops, omental thickening or ascites. Perfusion parameters (PEI, BF and BV) of the pancreatic lesions were (**Figure 5**) significantly reduced compared to the normal enhancing pancreas (**Table 2**). In view of encasement of the celiac axis by the pancreatic lesion, a ultrasound guided percutaneous FNAC was advised to confirm the nature of the lesion. FNAC revealed multiple necrotizing epithelioid cell granulomas with lymphocytic infiltration. A diagnosis of pancreatic tuberculosis was made and the patient was treated with ATT. On follow-up, significant improvement was seen both clinically and on imaging (ultrasound revealed decrease in size of pancreatic lesions).

Discussion

Isolated pancreatic involvement in tuberculosis is extremely rare probably because of the protective effect of pancreatic enzymes against mycobacteria which interferes with their seeding.^{1,2} Pancreatic TB is commonly seen in young patients, with female predominance, and past history of TB or in TB endemic regions.³ The first line non-invasive imaging modalities for pancreatic pathology include ultrasound and CT. Ultrasound may reveal focal hypoechoic or hypo- to isoechoic lesion or cystic collection most often in the body of pancreas followed by head and tail.² CT may show hypodense or cystic lesions in the pancreas, with features of acute or chronic pancreatitis and necrotic retroperitoneal or mesenteric lymphadenopathy along with bowel wall thickening and ascites. One of our patients presented with a hypodense mass in pancreatic head mimicking pancreatic carcinoma while the other had hypodense lesions in distal body and tail of pancreas with adjacent fascial thickening mimicking pancreatitis.

Volume perfusion CT is a new and upcoming modality which can quantify perfusion at microvascular level by measuring temporal changes in attenuation. VPCT was performed in both

these cases and it showed reduction of blood flow and blood volume in the lesions as compared to normal pancreas (**Tables 1 and 2**). Perfusion parameters are also reduced in pancreatic cancer however the extent of reduction is much more than in a pancreatic tubercular mass. To the best of our knowledge perfusion CT findings in pancreatic tuberculosis have not been published in literature. VPCT parameters may enable differentiation between pancreatic cancer and pancreatic tuberculosis, although further studies are needed to establish the statistical significance of differences in perfusion parameters.

AJAY KUMAR YADAV¹
JEYASEELAN NADARAJAH¹
DEVASENATHIPATHY KANDASAMY¹
RAJU SHARMA¹
SHIVANAND GAMANAGATTI¹
PRAMOD KUMAR GARG²

Correspondence: Dr Raju Sharma
Departments of Radiodiagnosis¹ and Gastroenterology,²
All India Institute of Medical Sciences,
New Delhi - 110029, India
Email: raju152@yahoo.com

References

1. Khaniya S, Koirala R, Shakya VC, Adhikary S, Regmi R, Pandey SR, et al. Isolated pancreatic tuberculosis mimicking as carcinoma: a case report and review of the literature. *Cases J*. 2010;**3**:18.
2. Nagar AM, Raut AA, Morani AC, Sanghvi DA, Desai CS, Thapar VB. Pancreatic tuberculosis: a clinical and imaging review of 32 cases. *J Comput Assist Tomogr*. 2009;**33**:136–41.
3. Xia F, Poon RT, Wang SG, Bie P, Huang XQ, Dong JH. Tuberculosis of pancreas and peripancreatic lymph nodes in immunocompetent patients: experience from China. *World J Gastroenterol*. 2003;**9**:1361–4.

Colonic spirochetosis in an immunocompetent child

Introduction

This is the first case report of colonic spirochetosis from a tropical country like India. Clinicians should consider rare

diagnosis like colonic spirochetosis if chronic dysentery is not responding to common medical treatment. Colonoscopy with multiple biopsies and a good histopathologic examination by an experienced pathologist should be done. Prognosis is favorable with medication. The clinical significance of intestinal spirochetes in humans remains highly debated. It remains unclear whether the presence of microorganisms coating the intestinal mucosa signifies mere colonization or disease.

Case report

A 10-year-old boy presented to our Gastroenterology outpatient department with the complaints of blood-mixed loose stools since last three months. He reported up to 10 bowel movements per day associated with lower abdominal pain and cramping that was relieved by defecation. He had lost 5 kg weight over last three months. He was treated at another hospital with various combinations of antibiotics, probiotics and anti-diarrheal drugs without any improvement in symptoms. On physical examination, the patient was in no acute distress. He was afebrile with a pulse rate of 68 per minute and respiratory rate of 17 per minute. Cardiac, abdominal, respiratory and nervous systems were unremarkable. An initial laboratory workup included complete blood count, stool for routine-microscopic examination and stool culture. His hemoglobin was 9 g/dl and his stool was positive for occult blood. However, no ova, cyst or parasite was detected in the stool. Stool culture was negative. Colonoscopy was done which

Figure 1: Colonoscopy showing erythematous, edematous and elevated lesion in the hepatic flexure

revealed three erythematous, edematous, elevated lesions with small superficial ulcers (2 to 3 mm) and normal intervening mucosa in the hepatic flexure and ascending colon (**Figure 1**). Histopathology by hematoxylin and eosin stain of the colonic biopsy revealed a fuzzy band on the surface of the enterocytes. This finding was highly suggestive of intestinal spirochetosis. These findings were further confirmed by Warthin-Starry staining, which revealed a moderate number of spirochetes in the colonic biopsies (**Figure 2**). Biopsies from esophagus, stomach and duodenum showed no pathology. Diagnosis of intestinal spirochetosis was made and the patient was put on oral metronidazole initially for 14 days. However, mild symptoms persisted even after 14 days and a repeat course of

Figure 2: Colonic spirochetosis a) hematoxylin and eosin stain; and b) Warthin-Starry stain

metronidazole was given for another 14 days. By the end of second course the patient improved clinically. Repeat colonoscopy after completion of therapy showed complete resolution of colonic ulcers. The patient was doing well till 12 weeks post-treatment follow-up.

Discussion

Adhesion of spirochetes to the mucosal brush border, resulting in its thickening as seen on light microscopy has been reported worldwide in both children and adults.¹ *Brachyspira aalborgi* and *Brachyspira pilosicoli* are members of *Brachyspiraceae* family. Both have been described in humans and are considered as cause of human intestinal spirochetosis.² Since 1997, *Brachyspira* spp. has been included in the list of human enteropathogenic bacteria.³ Both these spirochetes are slow growing fastidious anaerobes and require specific media, with estimated growth times of 6 days for *B. pilosicoli* and up to 2 weeks for *B. aalborgi*. Morphologically, they are coiled Gram-negative bacilli and are motile in liquid environment due to presence of flagellae. Generally they are considered non-invasive but systemic spread of *B. pilosicoli* has been documented in critically ill patients.⁴

These spirochetes are susceptible to several antibiotics, such as metronidazole, meropenem, chloramphenicol, ceftriaxone and tetracycline, but 60% isolates are resistant to ciprofloxacin.⁵ Mode of transmission is unclear, but is likely to occur by feco-oral route (contaminated water, colonized/infected feces). Due to higher prevalence in homosexual men, sexual transmission has also been suggested.⁶ Prevalence varies considerably across different geographic and immune conditions. In developed countries it ranges between 1.1–5.0%, with higher prevalence in homosexual men and HIV positive patients.¹ In most cases, intestinal spirochetosis is asymptomatic and presents incidentally during a screening colonoscopy for other reasons. However, infected children usually complain of persistent diarrhea, rectal bleeding, constipation, abdominal pain, weight loss, failure to thrive, nausea and lack of appetite.⁵ The severity of disease can vary from asymptomatic colonization to invasive and rapidly fatal progression, but there appears to be no correlation between degree of immunodeficiency in HIV positive patients and the extent of disease.¹ Colonic involvement is documented from distal to proximal, including rectum and appendix.⁶

Mucosal appearance on endoscopy is insufficient for

diagnosing infection, since it can present as normal, polypoid, and/or erythematous mucosa, or merely as non-specific lesions.⁷ The histologic appearance of spirochetes mimics a 'false brush border' formed by a layer of spirochetes. The surrounding cytostructure may show inflammation with slight oedema, infiltrate of monocytes, lymphocytes, plasma cells and neutrophils in the lamina propria, as well as elongated and hyperplastic crypts. On electron microscopy, spirochetes are attached perpendicularly to the epithelial membrane of the enterocytes and the microvilli appear shortened or depleted. Histologically it is not possible to distinguish *B. pilosicoli* from *B. aalborgi*. Genetic methods have been developed to identify *Brachyspira* species from stool and tissue samples.

Treatment strategies have been proposed for eradicating intestinal spirochetosis, including macrolides and clindamycin. But metronidazole seems to be the drug of choice, with a regimen of 500 mg 3 times a day for 10 days in adults and 15 mg per kg bodyweight 3 times per day for 5 days in children.⁵ However, longer therapy may be required sometimes as in the present case where it took 28 days for complete recovery. Currently, there is little evidence for the most effective antibiotic agent as treatment response is variable and sometimes ineffective, supporting the hypothesis that these microorganisms are harmless commensals.¹ Spontaneous recovery has been described after a prolonged period for up to 8 months.⁸

JAYANT KUMAR GHOSH
SUNDEEP KUMAR GOYAL
SANGEY CHOPEL LAMTHA
PANKAJ KAUSHIK
MANAS KUMAR BEHERA
ABHILASH VB
VINOD KUMAR DIXIT
AK JAIN

Correspondence: Dr. Jayant Kumar Ghosh
Department of Gastroenterology, Institute of Medical Sciences,
Banaras Hindu University, Varanasi, UP- 221005, India
Email: drjayantkg@yahoo.co.uk

References

1. Tsinganou E, Gebbers JO. Human intestinal spirochetosis—a review. *Ger Med Sci*. 2010;**8**:Doc01.
2. Paster BJ, Dewhirst FE. Phylogenetic foundation of spirochetes. *J Mol Microbiol Biotechnol*. 2000;**2**:341–4.
3. Bruckner DA, Colonna P. Nomenclature for aerobic and facultative bacteria. *Clin Infect Dis*. 1997;**25**:1–10.

4. Bait-Merabet L, Thille A, Legrand P, Brun-Buisson C, Cattoir V. Brachyspira pilosicoli bloodstream infections: case report and review of the literature. *Ann Clin Microbiol Antimicrob.* 2008;**7**:19.
5. Brooke CJ, Hampson DJ, Riley TV. In vitro antimicrobial susceptibility of Brachyspira pilosicoli isolates from humans. *Antimicrob Agents Chemother.* 2003;**47**:2354–7.
6. Helbling R, Osterheld MC, Vaudaux B, Jaton K, Nydegger A. Intestinal spirochetosis mimicking inflammatory bowel disease in children. *BMC Pediatr.* 2012;**12**:163.
7. Alsaigh N, Fogt F. Intestinal spirochetosis: clinicopathological features with review of the literature. *Colorectal Dis.* 2002;**4**:97–100.
- 8.. Marthinsen L, Willen R, Carlen B, Lindberg E, Varendh G. Intestinal spirochetosis in eight pediatric patients from Southern Sweden. *APMIS.* 2002;**110**:571–9.

Immunoproliferative small intestinal disease: a report of 6 cases

Introduction

Immunoproliferative small intestinal disease (IPSID) is a rare low-grade B-cell lymphoma arising from mucosa associated lymphoid tissue (MALT), representing approximately one-third of intestinal lymphoma. Histopathologically it is characterized by infiltration of small bowel wall with lymphocytes and plasma cells. The infiltrates produces an aberrant immunoglobulin (IgA), a truncated alpha heavy chain without the light chain component.^{1,2} Galian et al³ described the disease in three stages depending upon type of cellular infiltrate and mesenteric nodal involvement (**Table 1**). Presentation may mimic that of celiac disease, tropical sprue or parasitic infestation and patients are being treated on the line of celiac disease and other causes of malabsorption for prolonged duration before being diagnosed as IPSID/NHL.^{4,5} Emaciation and diarrhoea is due to malabsorption syndrome and protein losing enteropathy secondary to lymphoma infiltration of intestine. Most patients present during early stages and are curable with oral antimicrobials.⁶ Few may progress to high-grade lymphoma requiring systemic chemotherapy. We present here the clinico-

Table 1: The Galian staging system³

Stage	Small intestine	Lymph nodes
A	Lymphoplasmacytic or plasmacytic infiltration of the lamina propria, variable villous atrophy	Plasmacytic infiltration, nodal architecture generally preserved
B	Atypical lymphoplasmacytes or plasmacytes with immunoblast-like cells with extension to at least submucosa, subtotal or total villous atrophy	Atypical plasmacytic infiltrate with immunoblast-like cells, subtotal or total effacement of nodal architecture
C	Frankly malignant invasion through entire intestinal wall	Malignant effacement of entire lymph node

pathological characteristics, treatment and outcome of six patients treated at our institute during the last 10 years, along with a review of literature.

Case reports

A total of six patients with median age of 27 years (range: 15–35) were identified and treated. Median duration of symptom was 13.5 months (range: 6–26). All patients presented with history of recurrent watery loose motion with decreased appetite, weight loss and low grade fever. The median time from diagnosis of malabsorption to diagnosis of IPSID was 10 months (range: 1–20). Test for alpha heavy chain protein (AHCP) in serum was not done.

Barium meal follow through showed ileo-ileal intussusception in case number 4 and 6 (**Table 2**). All patients showed evidence of oedematous mucosal thickening in small bowel and enlarged mesenteric lymph nodes on CT scan. Endoscopic biopsy of mucosal lesions showed diffuse infiltrate of small atypical lymphoid cells or lymphoplasmacytic cells (**Figure 1A**) which were CD20⁺CD3⁺ with partial or total villous atrophy. Repeat biopsy during progression showed features of high grade B-cell lymphoma in case number 3 (**Figures 1B & 1C**).

Galian stage, treatment details and outcome are summarised in **Table 3**. Case number 1 received 6 cycles of CHOP (cyclophosphamide, doxorubicin, vincristine, and prednisolone) and achieved complete response (CR) lasting for 10 months and then was lost to follow-up. Case 2 also received 6 cycles of CHOP with tetracycline (1g/day) and was in stable condition for 1 year after which the disease progressed, and the patient succumbed 45 months after initial